

# **WI-FI ELECTROMAGNETIC FIELDS EXERT GENDER RELATED ALTERATIONS ON EEG**

**ARGIRO E. MAGANIOTI<sup>1</sup>, CHARALABOS C. PAPAGEORGIOU<sup>2,3</sup>,  
CHRISANTHI D. HOUNTALA<sup>1</sup>, MILTIADES A. KYPRIANOU<sup>3</sup>,  
ANDREAS D. RABAVILAS<sup>3</sup>, GEORGE N. PAPADIMITRIOU<sup>2</sup>,  
CHRISTOS N. CAPSALIS<sup>1</sup>**

*<sup>1</sup> NATIONAL TECHNICAL UNIVERSITY OF ATHENS, DEPARTMENT OF ELECTRICAL ENGINEERING, DIVISION OF INFORMATION TRANSMISSION SYSTEMS AND MATERIAL TECHNOLOGY, 9 Iroon Polytecneioy str., Athens, 15773, GREECE, Tel: ++30210-7722574, Fax: ++30210-7723520*

*<sup>2</sup> UNIVERSITY OF ATHENS,<sup>1ST</sup> DEPARTMENT OF PSYCHIATRY, EGINATION HOSPITAL, 74 Vas.Sophias Ave., Athens, 11528, GREECE*

*<sup>3</sup> UNIVERSITY MENTAL HEALTH RESEARCH INSTITUTE (UMHRI), 2 Soranou tou Efesiou (PO Box 66 517) Athens, 15601, GREECE, Tel: ++30210-6536902, Fax: ++30210-6537273*

## ***Abstract***

The present study investigated the influence of electromagnetic fields, similar to that emitted by Wi-Fi system, on brain activity. Fifteen female and fifteen male subjects performed a short memory task (Wechsler test), both without and with exposure to a 2.4GHz Wi-Fi signal. For each subject, radiation condition and electrode, the amplitude in the frequency domain of the EEG signal was calculated from the recordings of 30 scalp electrodes, using the Fourier transform.

The presence of radiation had no effect on the energies of alpha and beta band of male subjects, while it reduced these energies of female subjects, resulting in significantly lower energies, as compared to those of males. Delta and theta band energies did not experience any noteworthy effect from gender, radiation condition and their interaction. Conversely, there was a significant interaction effect (gender x radiation) on the energies of alpha and beta rhythms.

Interestingly, this pattern was observed for a number of electrodes, which formed two distinct clusters: one located at right- anterior and the second at occipital brain areas.

The present data support the idea that Wi-Fi signal may influence normal physiology through changes in gender related cortical excitability, as reflected by alpha and beta EEG frequencies.

## 1. INTRODUCTION

A lot of research has been done in the last years on the possible effects of Radio Frequency Electromagnetic Fields (RF - EMF) on biological matter. The majority of this research regards the potential health impacts of cell phones and other mobile communication emitters.

There are several studies involving subjects who perform various tasks while exposed to RF [1-6], which conclude that some aspects of cognitive function and some measures of brain physiology may be affected by the exposure to RF of the type emitted by cellular phones. Interestingly, the RF effect was found to be gender related [7, 8].

Based on the reviewed publications examining possible biological effects of RF exposure, the evidence suggests that the exposure to RF affect the human brain and its subsequent output in the form of cognition and behavior. This assumption is in line with recent reviews by Cook et al., (2006) and Valentini et al., (2007) [9, 10]. However, there are also reports contradicting this assumption Kleinlogel et al., (2008) [11].

Very common and constantly increasing sources of RF exposure are wireless networks that allow high-speed internet access and services, such as Wi-Fi. Inevitably, there has been concern about possible health effects from such exposure; however little research has been devoted to investigate the possible effects of Wi-Fi signal on biological systems.

Most of the studies that have been conducted, investigating the biological effects of Wi-Fi on humans, are mainly dealing with the amount of energy absorbed in human tissue and therefore limited in the measurement of the Specific Absorption Rate (SAR). There are also dosimetry studies ongoing, measuring RF levels around the globe, including that coming from wireless local area networks (WLANs) [12] and others which indicate that the exposure level is low compared to other sources [13]. Another approach, involved experimental procedures for whole-body RF exposure of animals in order to investigate the possible effects of Wi-Fi signal on biological systems [14].

In view of the above considerations, it can be hypothesized that the electrophysiological brain activity, as reflected by Electroencephalography (EEG) – alpha, beta, theta, delta bands – in association with cognitive task operations, could be of value in identifying possible pathophysiological alterations evoked by Wi-Fi signals and their connection with gender. Thus, the present study was designed to determine whether the presence of Wi-Fi signal affects the patterns of EEG activity elicited during a short memory task (Wechsler test).

## 2. METHODS

### 2.1. Participants

Thirty healthy individuals (15 men and 15 women, mean age =  $23.76 \pm 1.67$  years, mean education =  $16.9 \pm 1.06$  years) participated in the experiment. The participants were homogeneous with regards to age and educational level and had no history of any hearing problem. Informed consent was obtained from all subjects.


### 2.2. Experimental setup and Measurement Procedure

The subjects were evaluated with the digit span Wechsler Auditory test. A warning stimulus of either high (3000 Hz) or low frequency (500 Hz) was presented through earphones to the subjects, who were asked to memorize the numbers that followed. The warning stimulus lasted 100msec. A one second interval followed the onset of the warning stimulus and then the numbers to be memorized were presented by a female voice. At the end of the number sequence presentation, the same signal tone was repeated. The signals were recorded for a 1500msec interval, which means 500msec before the warning stimulus (EEG) and 1000msec after that (ERP), as described in previous articles [8, 15]. The numbers were recalled by the subject in the same (low frequency tone) or in the opposite order (high frequency tone) than that presented to the participant. The total task consisted of 52 repetitions for a period of about 45min. The subjects performed the tasks twice, with and without radiation, with an interval of two weeks between the measurements. The order in which the subject was exposed at the EMF (exposure at the first or second visit) was random. The EMF was emitted by a Wi-Fi access point that was operating at 2.4 GHz frequency. The Wi-Fi signal was radiated by a dual dipole antenna, with 20dBm power and

# WI-FI EXERT GENDER RELATED ALTERATIONS ON EEG

OFDM modulation. The access point was placed at a distance of 1.5m from the right part of their head. The field strength was 0.49V/m at the point where the subjects' head was standing. According to E. Kapareliotis et al. [16] there is no evidence that a Wi-Fi signal causes interference at the EEG recordings at the distance of 1.5m from the EEG electrodes.

The experiment was conducted in a Faraday room, which screened any electromagnetic interference that could affect the measurements. The attenuation of the mean field was more than 30 dB.



**Figure1.** Experimental Setup

The electrophysiological signals were recorded with Ag/AgCl electrodes. Electrode resistance was kept constantly below 5 k $\Omega$ . EEG activity was recorded from 30 scalp electrodes (FC6, FC2 F8, F4, Fz, AFz, Fp2, Fz, Fp1, P3, T3, FC1, FC5, F3, F7, T6, P4, CP6, CP2, T4, C4, O2, Oz, O1, Pz, Cz, T5, CP1, CP5, C3) based on the International 10-20 system of Electroencephalography [17], referred to both earlobes. Linked ear lobes served as reference. The bandwidth of the amplifiers was set at 0.05 – 35 Hz. Eye movements were recorded with the use of electro-oculogram (EOG) and recordings with EEG higher than 75  $\mu$ V were excluded. The evoked bio potential signal was submitted to an analogue-to-digital conversion, at a sampling rate of 1 KHz and was averaged by a computerized system.

## 2.3. Data Transformation

For each question 1500 data points, each corresponding to time segments of 1msec duration for each electrode were saved. In order to optimize the signal to noise ratio for each subject and each channel all EEG values were average referenced on the basis of the grand average across the 52 repetitions of the EEG values. This procedure was done separately for each EMF condition. The final data for analysis for each subject and condition consisted of 1500 amplitude values for each electrode, expressed in  $\mu$ Volts corresponding to the 1500msec of the time period [8, 15], 500ms before the onset of the first warning stimulus (EEG), and 1000 ms after the onset (ERP).


For each subject, each radiation condition and each electrode, the amplitude in the frequency domain was calculated using the Fourier transform (FFT) (EEG). This analysis confirmed the expected pattern of the spectral distribution of the signals, with the appearance of peaks within the range of the four basic rhythms d (0–4 Hz),  $\theta$  (5–7 Hz),  $\alpha$  (8–13 Hz) and  $\beta$  (14–30 Hz) in all EEG series.

## 2.4. Statistical analysis

The EEG energies were log-normalized so that their distribution for all the electrodes and both experimental conditions did not deviate from normality. For each band, the energies at the thirty electrodes were subjected to multivariate analysis of variance with gender (male-female) and radiation condition (on-off) as the independent factors, followed by post-hoc pair wise comparisons with Bonferroni corrections. The significance level was set at 0.05.

## 3. RESULTS


Multivariate analysis of variance did not reveal any significant effect of gender, radiation condition and their interaction on the energies of the delta and theta bands. Conversely, there was a significant interaction effect (gender x radiation) on the energies of the alpha and beta band. The nature of this interaction effect can be clarified by Figure 2, which shows the average values at the alpha band for male and female subjects, for the two radiation conditions at electrode F8. As this figure shows, at the absence of EMF the alpha band energies of the two genders are practically the same. The presence of radiation did not have any effect on the energy of male subjects, whereas it reduced the energy of female subjects, resulting in a significantly lower energy, as compared to the energy of male subjects.



**Figure 2.** Average values at the alpha band for male and female subjects, for the two radiation conditions at electrode F8

The same pattern was observed for a number of electrodes (FC6, F8, Fp2, Fpz, C4, O2, Oz and O1). As Figure 3 shows these electrodes form two distinct clusters, one located at right- anterior and the second at occipital brain areas. Interestingly, analysis revealed that the beta band energies have practically the same behavior.

## WI-FI EXERT GENDER RELATED ALTERATIONS ON EEG



**Figure 3.** p-values of the differences between male and female subjects in the absence and presence of the Wi-Fi signal

### 3. SUMMARY - DISCUSSION

The comparison between experimental conditions showed that the presence of radiation did not have any effect on the energies of the alpha and beta band of male subjects, while it reduced these energies of female subjects, resulting in significantly lower energies, as compared to the energies of male subjects.

Additionally, the energies of the delta and theta bands did not experience any significant effect from gender, radiation condition and their interaction. Conversely, there was a significant interaction effect (gender x radiation) on the energies of the alpha and beta rhythms.

Interestingly, this pattern was observed for a number of electrodes, which formed two distinct clusters, one located at right-anterior and the second at occipital brain areas.

The results of the present study may be interpreted in the light of the psycho physiological and brain-imaging studies related to EEG functional anatomy. The quantification of EEG was proven a useful and appropriate method in measuring the level and topographical distribution of cortical activation during cognitive task performance. In general, the functional significance of varying brain activity can be seen in the vicinity to the underlying neural circuits. For instance, it is assumed that alpha band activity reflects an increased excitability level of neurons in the certain cortical areas, which may be related to an enhanced information transfer in thalamocortical circuits and is strongly correlated with working memory as well as with long-term memory engrams [18-21]. Beta bursts being related to cortico-cortical interactions shift the system to an attention state that consequently allows for gamma synchronization and perception [22-24].

The effect of Wi-Fi exposure (significant interaction effect -gender x radiation-on the energies of the alpha and beta bands) are in accordance with previous studies of our team regarding gender related differences in the EEG under EMF exposure of 900MHz and 1800MHz similar to that of mobile phones [8, 15, 25, 26, 5, 7]. Also Smythe and Costall (2003) [27] have reported sex-dependent effects of EMF exposure on human memory during a memory task.

Although, the biological basis for these sex differences remains elusive, emerging evidence provides plausible mechanisms for the explanation of these differences. In particular, central nervous system effects of EMFs have been considered to be secondary to damage to the blood-brain barrier (BBB) permeability [28-31]. At this point is reasonable to consider the existence of gender-related blood-barrier differences, a fact which would explain the fundamental differences between males and females in the intrinsic cognitive processes and in the way they are affected by different types of electromagnetic radiation [32-34]. Furthermore research also indicates that EMF exposure affects the melatonin release, specifically it has been demonstrated a reduced excretion of the urinary metabolite of melatonin among persons using a mobile phone for more than 25 min per day [35]. This observation would be better understood taking into account that in a study of pubertal individuals it has been

found significantly higher total, nocturnal and diurnal 6-sulfatoxymelatonin excretion in girls [36].

It is concluded that Wi-Fi may influence normal physiology through changes in gender related cortical excitability as it is reflected by the alpha and beta EEG frequencies.

#### 4. REFERENCES

- [1] D. Hamblin, A. Wood, R. Croft, C. Stough, Examining the effects of electromagnetic fields emitted by GSM mobile phones on human event-related potentials and performance during an auditory task, *Clinical Neurophysiology* 115 (2004) 171-178.
- [2] R. Jech, K. Sonka, E. Ruzicka, A. Nebuzelsky, J. Bohm, M. Juklickova, S. Nevsimalova, Electromagnetic field of mobile phones affects visual event related potential in patients with narcolepsy, *Bioelectromagnetics* 22 (2001) 519-528.
- [3] C. Krause, L. Sillanmaki, M. Koivisto, A. Haggqvist, C. Saarela, A. Revonsuo, M. Laine, H. Hamalainen, Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task, *Neuroreport* 11 (2000) 761-764.
- [4] E. Maby, R. Le Bouquin Jeannes, G. Faucon, Short-term effects of GSM mobile phones on spectral components of the human electroencephalogram, *Proceedings of Engineering in Medicine and Biology Society* (2006) 3751-3754.
- [5] A. Maganioti, C. Hountala, C. Papageorgiou, A. Rabavilas, G. Papadimitriou, C. Capsalis, Cointegration of ERP signals in experiments with different EMF conditions, *Health* 2 (2010) 400-406.
- [6] C. Papageorgiou, E. Nanou, V. Tsiafakis, E. Kapareliotis, K. Kontoangelos, C. Capsalis, A. Rabavilas, C. Soldatos, Acute mobile phone effects on pre-attentive operation, *Neuroscience Letters* 397 (2006) 99-103.
- [7] A. Maganioti, C. Hountala, C. Papageorgiou, M. Kyprianou, A. Rabavilas, C. Capsalis, Principal component analysis of the p600 waveform: RF and gender effects, *Neuroscience Letters* 478 (2010) 19-23.
- [8] C. Papageorgiou, E. Nanou, V. Tsiafakis, C. Capsalis, A. Rabavilas, Gender related differences on the EEG during a simulated mobile phone signal, *Neuroreport* 15 (2004) 2557-2560.
- [9] C. Cook, D. Saucier, A. Thomas, F. Prato, Exposure to ELF magnetic and ELF-modulated radiofrequency fields: the time course of physiological and cognitive effects observed in recent studies, *Bioelectromagnetics* 27 (2006) 613-627.
- [10] E. Valentini, G. Cursio, F. Moroni, M. Ferrara, L. De Gennaro, M. Bertini, Neurophysiological effects of mobile phone electromagnetic fields on humans: a comprehensive review, *Bioelectromagnetics* 28 (2007) 415-432.
- [11] H. Kleinlogel, T. Dierks, T. Koenig, H. Lehmann, A. Minder, R. Berz, Effects of weak mobile phone - electromagnetic fields (GSM, UMTS) on event related potentials and cognitive functions, *Bioelectromagnetics* 6 (2008) 488-97.
- [12] K. Foster, Radiofrequency Exposure from Wireless LANS Utilizing Wi-Fi Technology, *Health Physics* 92 (2007) 280-289.
- [13] M. Martínez-Búrdalo, A. Martín, A. Sanchis, R. Villar, FDTD assessment of human exposure to electromagnetic fields from Wi-Fi and bluetooth devices in some operating situations. *Bioelectromagnetics* 30 (2009) 142-151.
- [14] C. Marino, P. Galloni, F. Nasta, R. Pinto, C. Pioli, G. Lovisolo, Measures for the exposure of newborn animals to Wi-Fi signals, *Proceedings of Open questions in the research on biological and health effects of low-intensity RF-EMF* (2008), 12.
- [15] E. Nanou, V. Tsiafakis, E. Kapareliotis, C. Papageorgiou, A. Rabavilas, C. Capsalis, Influence of the interaction of the 900 MHz signal with gender on EEG energy: experimental study on the influence of 900 MHz

## WI-FI EXERT GENDER RELATED ALTERATIONS ON EEG

radiation on EEG, *Environmentalist* 25, (2005) 173-179.

[16] E. Kapareliotis, E. Nanou, V. Tsiafakis, A. Sotiriou, L. Pragiatis, C. Capsalis, Electromagnetic compatibility between Wi-Fi access point and EEG signals, *Proceedings of 4<sup>th</sup> International Workshop Biological Effects of Electromagnetic Fields* (2006) 545-551, Crete-Greece.

[17] H. Jasper, The ten-twenty electrode system of the international federation, *Electroencephalography Clinical Neurophysiology* 10 (1958) 371–375.

[18] C. Neuper, G. Pfurtscheller, Event-related dynamics of cortical rhythms: frequency-specific features and functional correlates, *International Journal of Psychophysiology* 43 (2001) 41-58.

[19] O. Jensen, J. Gelfand, J. Kounios, J. Lisman, Oscillations in the alpha band (9-12 Hz) increase with memory load during retention in a short-term memory task, *Cerebral Cortex*. 12 (2002) 877-882.

[20] S. Leiberg, W. Lutzenberger, J. Kaiser, Effects of memory load on cortical oscillatory activity during auditory pattern working memory. *Brain Research* 1120 (2006) 131-140.

[21] S. Palva, M. Palva, New vistas for alpha-frequency band oscillations. *Trends in Neurosciences* 30 (2007) 150-158.

[22] A. Wrobel, Beta activity: a carrier for visual attention, *Acta neurobiologiae experimentalis (Wars)* 60 (2000) 247-260.

[23] R. Gaillard, S. Dehaene, C. Adam, S. Clémenceau, D. Hasboun, M. Baulac, L. Cohen, L. Naccache, Converging intracranial markers of conscious access, *PLoS Biology* 7 (2009) e61.

[24] S. Pockett, G. Bold, W. Freeman, EEG synchrony during a perceptual-cognitive task: widespread phase synchrony at all frequencies, *Clinical Neurophysiology* 120 (2009) 695-708.

[25] C. Hountala, A. Maganioti, C. Papageorgiou, E. Nanou, M. Kyprianou, V. Tsiafakis, A. Rabavilas, C. Capsalis, The spectral power coherence of the EEG under different EMF conditions. *Neuroscience Letters* 441 (2008) 188-192.

[26] E. Nanou C. Hountala, A. Maganioti, C. Papageorgiou, V. Tsiafakis, A. Rabavilas, C. Capsalis, Influence of a 1,800 MHz electromagnetic field on the EEG energy, *Environmentalist* 29 (2009) 205-209.

[27] J. Smythe, B. Costall, Mobile phone use facilitates memory in male, but not female, subjects, *Neuroreport* 14 (2003) 243-246.

[28] L. Salford, et al., Permeability of the blood brain barrier induced by 915MHz electromagnetic radiation continuous wave and modulated at 8, 16, 50 and 200 Hz, *Microscopy Research and Technique* 27 (1994) 535–542.

[29] A. Schirmacher, Electromagnetic fields (1.8 GHz) increase the permeability of sucrose of the blood–brain barrier in vitro, *Bioelectromagnetics* 21 (2000) 338–345.

[30] L. Salford, A. Brun, et al., Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones, *Environmental Health Perspectives* 111 (2003) 881–883.

[31] J. Eberhardt, B. Persson, A. Brun, L. Salford, L. Malmgren, Blood–brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones, *Electromagnetic Biology and Medicine* 27 (2008) 215–229.

[32] W. Skrandies, P. Reik, C. Kunze, Topography of evoked brain activity during mental arithmetic and language tasks: sex differences, *Neuropsychologia* 37 (1999), 421-430.

[33] W. Dimpfel, W. Wedekind, I. Keplinger, Gender difference in electrical brain activity during presentation of various film excerpts with different emotional content, *European Journal of Medical Research* 8 (2003): 192-198.

- [34] M. Briere, G. Forest, S. Chouinard, R. Godbout, Evening and morning EEG differences between young men and women adults, *Brain and Cognition* 53 (2003) 145-148.
- [35] J. Burch, J. Reif, C. Noonan, T. Ichinose, A. Bachand, T.. Koleber, M. Yost, Melatonin metabolite excretion among cellartelephone users, *International Journal of Radiation Biology* 78 (2002) 1029–1036.
- [36] H. Fideleff, H. Boquete, G. Fideleff, L. Albornoz, S. Lloret, M. Suarez, A. Esquifino, M. Honfi, D. Cardinali, Gender-related differences in urinary 6-sulfatoxymelatonin levels in obese pubertal individuals, *Journal of pineal research* 40 (2006) 214-218.